REDISCOVERY OF THE SOUTH INDIAN POPULATION OF THE BRUSH FLITTER BUTTERFLY HYAROTIS MICROSTICTUM COORGA FROM KARNATAKA

PRADEEP NAYAK

'Sannidhi',#12-104/4, Kongoor Mutt Road, Kulshekar, Mangalore - 575 005

Email: nayakpradeep007@gmail.com

Reviewer: Peter Smetacek

INTRODUCTION

The southern Indian population of the Brush Flitter Hyarotis microstictum coorga Evans, 1949 is a skipper butterfly whose known distribution is restricted to Karnataka. The butterfly is so rare that there is no live image available and the species has not been sighted in the last 95 vears, the most recent confirmed record of the species being from 1927 (Yates, 1931). However, during our recent surveys in Sampaje Ghats of Karnataka we regularly observed the butterfly from 18th March to 18th April 2022. Our observations suggest that the butterfly is restricted to small pockets of Sampaje Ghats near Payaswini at an elevation of 225 m. We surveyed all the localities around it and found that the butterfly was found on both sides of the road and individuals were found nectaring on the roadside flowers of Leea indica. It is also seen at similar elevation in NE India viz. 20 m-200 m (Monsoon Jvoti Gogoi, 2022 pers. comm..).

The taxon *coorga* was originally described as a subspecies of *H. microstictum* by Evans (1949). *H. microstictum* has a distribution range from Assam to Myanmar, Thailand, Vietnam, Langkawi, W. Malaysia, Borneo, Sumatra, Luzon (Y. Inayoshi, 1996-2022). Although *Hyarotis coorga* is treated as a species in some web

resources, the change in status from subspecies to species is not validated in any published literature and appears to be frivolous. Evans (1949) and Gaonkar (1996) treated it as subspecies of *H. microstictum*. The taxon was described on the basis of a male collected from Sati river, Coorg by J.A. Yates on 2nd December 1927. The typical subspecies on the other hand was described by Wood-Mason & de Nicéville (1887) from Cachar hills of Assam.

The subspecies *coorga* differs from *H. m. microstictum* in having larger white spots on the upperside forewing (Evans, 1949). *H. m. coorga* is rare in Karnataka (Gaonkar, 1996) and *H. m. microstictum* is very rare and variable according to Evans (Yates, 1931).

DISCUSSION

Our photographs of *H. m. coorga* are the first ever of the taxon in nature. Our observations suggest that the species is locally not rare in Sampaje Ghats, although the species is reportedly very rare (Yates 1931). The species was seen at an elevation of around 200 m and the recent records from *H. m. microstictum* from S. Cachar hills and Jaintia hills in NE India also share the same elevation range with *H. m. coorga*.



Hyarotis microstictum coorga, April, Sampaje Ghat of Karnataka ♂



Hyarotis microstictum coorga, April, Sampaje Ghat of Karnataka ♂



Hyarotis microstictum coorga, April, Sampaje Ghat of Karnataka ♀



Hyarotis microstictum coorga, April, Sampaje Ghat of Karnataka



Hyarotis microstictum (Feb 2012, Jaintia hills, Meghalaya, Pic: Monsoon J Gogoi)



Hyarotis microstictum (Feb 2012, Jaintia hills, Meghalaya, Pic: Monsoon J Gogoi)∂

ACKNOWLEDGEMENTS

I thank Isaac Kehimkar for his continuous encouragement. I also thank Monsoon J Gogoi for helping in preparing the manuscript and permission to use his photographs.

REFERENCES

Evans, W.H. 1949. A catalogue of the Hesperiidae from Europe, Asia & Australia in the British Museum (Natural History). The British Museum. London. 502 pp.

Wood-Mason, J. & L. de Nicéville. [1887]. :List of the lepidopterous insects collected

in Cachar by Mr. J. Wood-Mason. Part II. Rhopalocera. *J. Asiat. Soc. Bengal* 2. 55(4): 343-393, 4 pls.

Yates, J.A. 1931. The Butterflies of Coorg. *Journal of the Bombay Natural History Society* 34(4): 1003–1014.

Gaonkar, H. 1996. Butterflies of the Western Ghats, India (including Sri Lanka) - A Biodiversity Assessment of a Threatened Mountain System. A report submitted to the Centre for Ecological Sciences, Bangalore, India. 86 pp.