

***LILIUM LONGIFLORUM* AND *LILIUM BULBIFERUM* (LILIACAE):**

NEW LARVAL HOST PLANTS FOR *KANISKA CANACE* (LEPIDOPTERA: NYMPHALDAE) FROM ARUNACHAL PRADESH, INDIA

RUKSHA LIMBU¹ & ROSHAN UPADHAYA²

¹Department of Life Science, RIMT University, Mandi Gobindgarh, Punjab, 147301

²*Arunachal Pradesh Police, Miao, Arunachal Pradesh 792122

Corresponding author: roshanupadhaya11@gmail.com

Reviewer: Peter Smetacek

ABSTRACT

Lilium longiflorum and *Lilium bulbiferum* are reported as hitherto unrecorded larval host plants of *Kaniska canace* (Blue Admiral) from Arunachal Pradesh.

Key words: *Lilium longiflorum*, *Lilium bulbiferum*, Arunachal Pradesh

INTRODUCION

Kaniska Moore, 1899 is a monotypic genus of Nymphalidae, represented in India by two subspecies, *K. canace canace* (Linnaeus, 1763) distributed throughout the Himalayan region to N.E. India and *K. c. viridis* Evans, 1924, which is restricted to the Western Ghats (Varshney & Smetacek, 2015; Kehimkar, 2016). However, this species was also reported from the plain of Punjab (Singh et al., 2016) and Rajasthan (Sengupta, 2021). It is on the wing from March to December between an elevation of 1000m to 3000m in the Himalayan region and between 1,000m to 2,200m in

the southern hills of India (Kehimkar, 2016).

OBSERVATION

On 09.iv.2021, an adult female of *K. canace* was encountered laying six greenish spherical eggs on the lower surface of the leaves and stem of *Lilium longiflorum* and *Lilium bulbiferum* plants in a private garden of village Gaherigaon (27°11'53.9592" N; 96°57'49.6548" E) of Vijaynagar, district Changlang, Arunachal Pradesh. The eggs had longitudinal ribs and were laid singly.

The plants with the eggs were shifted to flower pots and placed inside the insectarium box made up of netting. The insectarium box was placed in the garden to rear the caterpillars under natural conditions. The caterpillars were monitored regularly and frass was cleaned every day. The different instars of the caterpillars were photographed using a

Canon DSLR camera with 90mm macro lens (figures 1-12).

All the six eggs hatched successfully on 16.iv.2021. The second instar caterpillars started feeding on leaves voraciously. On 16.v.2021 all the six larvae stop feeding and were observed moving away from the host plant to the bamboo branches which were kept in the insectarium to give support to the host plants; no frass was observed on that day. On 17.v.2021, all the six caterpillars were observed hanging vertically upside down on the bamboos from a silk pad and all the caterpillars pupated successfully between 12 pm to 1 pm. On 29.v.2021 all the pupae turned black and become wet. On 30.v.2021 adults of *Kaniska canace* emerged between 11am to 2 pm from all the six pupae. After expanding and drying their wings for 30 to 35 minutes each, the butterflies flew away.

Previously *Smilax* species have been reported as larval host plants of *K. canace* from India, notably *Smilax zeylanica* (Wynter-Blyth, 1957; Karmakar et al., 2018; Nitin et al., 2018). Robinson et al. (2001) recorded *Trycirtis hyrta* (Convallariaceae), *Lilium lancifolium* (Liliceae) and *Smilax china* (Smilacaceae) from China, and *Heterosmilax japonica*, *Smilax perfoliata*, *Smilax aspersa*, *S. arisanensis*, *S. aspericaulis*, *S. bracteata*, *S. lanceifolia*, and *S. riparia* (Smilacaceae) from other parts of the range of *K. canace*. Thus, the present study confirmed *Lilium longiflorum* and *Lilium bulbiferum* (Liliaceae) as previously unreported larval host plants for *K. canace*.

ACKNOWLEDGEMENT

The authors are thankful to Mrs. Dew Kumari Rai and Mr. Chandra Kumar Limbu for identifying the larval host plants.

REFERENCES

- Karmakar, T., N. Ravikantachari, V. Sarkar, S. Baidya, S. Mazumdar, V.K. Chandrasekharan, R.P. Das, Girish Kumar G.S., S.A. Lokhande, J. Veino, L. Veino, R. Veino, Z. Mirza, R. Sanap, B. Sarkar & K. Kunte. 2018. Early stages and larval host plants of some northeastern Indian butterflies. *Journal of Threatened Taxa*. 10(6): 11780-11799. [10.11609/jott.3169.10.6.11780-11799](https://doi.org/10.11609/jott.3169.10.6.11780-11799).
- Kehimkar, I. 2016. *Butterflies of India*. Bombay Natural History Society, Mumbai. 505 pp.
- Nitin, R., V.C. Balakrishnan, P.V. Churi, S. Kalesh, S. Prakash & K. Kunte. 2018. Larval host plants of the butterflies of the Western Ghats, India. *Journal of Threatened Taxa* 10(4): 11495–11550; <http://doi.org/10.11609/jott.3104.10.4.11495-11550>
- Robinson, G.S., P.R. Ackery, I.J. Kitching, G.W. Beccaloni & L.M. Hernández. 2001. Hostplants of the moth and butterfly caterpillars of the Oriental Region. The Natural History Museum, London and Southdene Sdn. Bht., Kuala Lumpur. 744 pp.
- Sengupta, D. 2021. First record of Blue Admiral *Kaniska canace* (Linnaeus, 1763)

(Lepidoptera: Nymphalidae) from the state of Rajasthan, India. *Revista Chilena de Entomología*. 47. 177-181. 10.35249/rche.47.1.21.17.

Singh, V., J.S. Kirti & D. Mehra. 2016. Butterflies of district Hoshiarpur, Punjab, India. *Indian Forester* 142 (10): 99- 104.

Varshney, R.K. & P. Smetacek. 2015. *A Synoptic Catalogue of the Butterflies of*

India. Butterfly Research Centre, Bhimtal and Indinov Publishing, New Delhi. ii + 261 pp., 8 pl.

Wynter-Blyth, M.A. 1957. *Butterflies of the Indian region*. Bombay Natural History Society, Bombay. xx + 523 pp., 72 pl.



Figure 1



Figure 2

Figure 1 & 2: *Lilium bulbiferum*

Figure 3 & 4: *Lilium longiflorum*Figure 5: Larval host plant
voraciously eaten by *Kaniska*
canace caterpillars

Figure 6: Egg hatching out



Figure 7



Figure 8



Figure 9



Figure 10



Figure 11




Figure 12

Figs. 7 -12: Early stages of *Kaniska canace*. 7: newly hatched larva. 8: larva resting after feeding. 9: Final instar. 10: larva pupating. 11: pupa. 12: adult ready to take flight